

b.) Amendment to the Claims

1. (Currently Amended) A method for producing a ~~neural crest~~ cell ~~expressing or~~ a neural ~~tube cell~~ surface marker, comprising:

culturing an embryonic stem cell *in vitro* in the absence of retinoic acid and in the presence of a stroma cell without forming embryoid body ~~for a time period from 1 day to 14 days~~, wherein the stroma cell is OP9 cell or PA6 cell ~~cells is recognized by a monoclonal antibody produced by hybridoma FERM BP 7573; and then~~

~~culturing continuously the cell in vitro in the absence of retinoic acid and in the presence of a stroma cell without forming embryoid body, wherein the stroma cell is recognized by a monoclonal antibody produced by hybridoma FERM BP 7573.~~

Claims 2-14 (Cancelled).

15. (Previously Presented) The method according to any one of claims 1, 80 or 81, which further comprises culturing under serum-free culture conditions.

Claims 16-17 (Cancelled)

18. (Previously Presented) The method according to any one of claims 1, 80 or 81, wherein the stroma cell is a stroma cell whose proliferation potency is deleted by a physicochemical treatment.

19. (Previously Presented) The method according to any one of claims 1, 80 or 81, wherein the stroma cell is a stroma cell whose proliferative potency is deleted by an antitumor agent, irradiation or pathologic tissue fixative.

20. (Previously Presented) The method according to claim 18, wherein the physicochemical treatment is an antitumor agent selected from the group consisting of mitomycin C, 5-fluorouracil, adriamycin and methotrexate.

21. (Previously Presented) The method according to any one of claims 1, 80 or 81, wherein the stroma cell is a stroma cell whose proliferative potency is deleted by a microwave fixation, a rapid freeze-substitution fixation, a glutaraldehyde fixation, a p-formaldehyde fixation, a formalin fixation, an acetone fixation, a Van fixation, a periodic acid fixation, a methanol fixation or an osmic acid fixation.

Claim 22 (Cancelled).

23. (Currently Amended) The method according to any one of claims 1, 80 or 81, wherein the stroma cell is ~~selected from the group consisting of: a fetal primary culture fibroblast; an SHM mouse derived STO cell; a mouse fetus derived NIH/3T3 cell; an M-CSF deficient mouse calvaria-derived OP9 cell; a mouse calvaria derived MC3T3-G2/PA6 cell; an embryonic stem cell derived stroma cell; and a bone marrow mesenchymal stem cell derived stroma cell.~~

24. (Previously Presented) The method according to any one of claims 1, 80 or 81, wherein the embryonic stem cell is selected from the group consisting of:

- (a) an embryonic stem cell established by culturing an early embryo before implantation;
- (b) an embryonic stem cell established by culturing an early embryo produced by nuclear transplantation of the nucleus of a somatic cell; and
- (c) an embryonic stem cell in which a gene on the chromosome of the embryonic stem cell of (a) or (b) is modified using gene engineering.

Claims 25-73 (Cancelled).

74. (Previously Presented) The method according to claim 23, wherein the embryonic stem cell is selected from the group consisting of:

(a) an embryonic stem cell established by culturing an early embryo before implantation;

(b) an embryonic stem cell established by culturing an early embryo produced by nuclear transplantation of the nucleus of a somatic cell; and

(c) an embryonic stem cell in which a gene on the chromosome of the embryonic stem cell of (a) or (b) is modified using gene engineering.

75. (Previously Presented) The method according to claim 74, wherein the stroma cell is recognized by a monoclonal antibody produced by hybridoma FERM BP-7573.

Claims 76-79 (Cancelled).

80. (Currently Amended) A method for producing a dopaminergic neuron, an acetylcholinergic neuron, a γ -aminobutyrate neuron or a serotonergic neuron, comprising:

culturing an embryonic stem cell *in vitro* in the absence of retinoic acid and in the presence of a stroma cell without forming embryoid body, wherein the stroma cell is ~~recognized by a monoclonal antibody produced by hybridoma FERM BP-7573~~ OP9 cell or PA6 cell.

81. (Currently Amended) A method for producing a neural stem cell which is stained by an anti-nestin antibody comprising:

culturing for a time period from 1 to 14 days an embryonic stem cell *in vitro* in the absence of both retinoic acid and BMP-4 and in the presence of a stroma cell without forming embryoid body, wherein the stroma cell is ~~recognized by a monoclonal antibody produced by hybridoma FERM BP-7573~~ OP9 cell or PA6 cell.

82. (Previously Presented) The production method according to claim 1, wherein the method produces said neural crest cell.

83. (Previously Presented) The production method according to claim 1, wherein the method produces said neural tube cell.

84. (Previously Presented) The production method according to claim 80, wherein the method produces said dopaminergic neuron.

85. (Previously Presented) The production method according to claim 80, wherein the method produces said acetylcholinergic neuron.

86. (Previously Presented) The production method according to claim 80, wherein the method produces said γ -aminobutyrate neuron.

87. (Previously Presented) The production method according to claim 80, wherein the method produces said serotonergic neuron.

88. (New) The method according to any one of claims 1, 80 or 81, wherein the stroma cell is a mouse calvaria-derived MC3T3-G2/PA6 cell.

89. (New) The method according to claim 23, wherein the stroma cell is recognized by a monoclonal antibody produced by hybridoma FERM BP-7573.

90. (New) The method according to claim 88, wherein the stroma cell is recognized by a monoclonal antibody produced by hybridoma FERM BP-7573.